

Classification  
 Order Actiniaria  
 Family Hormathiidae  
 Genus Hormatia  
 Species ***Hormathia digitata*** (Müller, 1776)

Reference: [Riemann-Zürneck K. 1994. Taxonomy and ecological aspects of the subarctic sea anemones \*Hormathia digitata\*, \*Hormathia nodosa\* and \*Allantactis parasitica\* \(Coelenterata, Actiniaria\). \*Ophelia\* 39\(3\): 197-224.](#)

Tissue	Cnidae type	Range Length x Width
Tentacles	basitrichs 1*	25-35 x 2.5-3
	basitrichs 2**	13-15.5 x 1.5-2
	spirocysts maximal	50 x 5-6
Pedal disc	basitrichs 1	19-23.5 x 3-3.5
	basitrichs 2	12-14 x 2
Column	basitrichs 1	16-19 x 3-3.5
	basitrichs 2	9-13.5 x 1.5-2
	p-mastigophore***	19-24 x 4-4.5
Actinopharynx	basitrichs 1	29-35 x 3-4
	basitrichs 2	14-17.5 x 1.5
	p-mastigophore	20.5-25 x 4-4.5
Mesenterial filaments	basitrichs 1	29-35 x 3-3.5
	basitrichs 2	14.5-17.5 x 1.5-2
	p-mastigophore	22-27 x 4-4.5
Acontia	basitrichs 1	29-36 x 3-3.5
	basitrichs 2	15.5-20 x 1.5-2

Notes: \*rare, \*\*very rare, \*\*\*in the scapulus, sometimes absent




Fig. 3. *Hormathia digitata*. Cnidome. The numbers refer to those with asterisks in Table 1. - P.DISC = Pedal disc, ACONT. = Acontia.

Source: Riemann-Zürneck (1994)

