

| | |
|---------|--|
| | Classification |
| Order | Actiniaria |
| Family | Hormathiidae |
| Genus | Phelliactis |
| Species | <i>Phelliactis lophohelia</i> Riemann-Zürneck, 1973 |

Reference: [Riemann-Zürneck K. 1973. Actiniaria des Südwatlantik. Hormathiidae, I. Helgolander wiss. Meeresunters 25: 273-325.](#)

| Tissue | Cnidae type | Range Length x Width* |
|-----------------------|--------------------|-----------------------|
| Tentacles | basitrichs 1 | 36.5-47 x 3-3.5 |
| | basitrichs 2 | 18** |
| | spirocysts maximal | 58-65 |
| Column | basitrichs 1 | 22.1-31 x 3-3.5 |
| | p-mastigophore | 34-41.5 x 5 |
| Actinopharynx | basitrichs 1 | 35-47 x 3-3.5 |
| | basitrichs 2 | 16** |
| | p-mastigophore | 31-40 x 4-4.5 |
| Mesenterial filaments | basitrichs 1 | 39-43 x 3 |
| | basitrichs 2 | 13-21 x 2-2.5 |
| | p-mastigophore | 26-36.5 x 4 |
| Acontia | basitrichs 1 | 36.5-48 x 3-3.5 |
| | basitrichs 2 | 18-24.5 x 2-2.5 |

Notes:

*All ranges were built taking maximal and minimal values obtained of two holotypes (68/10 and 68/13)

**some animals

No photo available